

Countability and Separation Axioms

Def. A space X is said to have a **countable basis** at $x \in X$ if there is a countable collection \mathcal{B} of neighbourhoods of x such that any neighbourhood of x contains an element of \mathcal{B} .

A space that has a countable basis at each of its points is said to be **first countable**.

First countable

For example, any metrizable space is first countable — consider $\{B_d(x, 1/n) : n \in \mathbb{N}\}$

Theo. Let X be a topological space.

- (4.1) a) Let $A \subseteq X$. If there is a sequence of points in A that converges to $x \in X$, then $x \in \bar{A}$. The converse holds if X is first countable.
- b) Let $f: X \rightarrow Y$ be continuous. If (x_n) be a sequence of points in X that converges to x , then $f(x_n) \rightarrow f(x)$. The converse holds if X is first countable.

Proof. Recall that we have proved the above for metrizable X by considering $B_d(x, 1/n)$.

Here, just consider

$$B_n = U_1 \cap \dots \cap U_n \text{ instead,}$$

where $\{U_n\}$ forms a countable basis at x .

The proof is relatively straightforward — let $x_n \in B_n \cap A$ for each n .

For (b), show that $f(\bar{A}) \subseteq \overline{f(A)}$. \square

Def. A topological space X is said to be **second countable** if it has a countable basis (for the topology).

Second countable

Prop. (4.2) Any second countable space is first countable.

The proof is direct.

For example, \mathbb{R} is second countable — consider $\{(a,b) : a,b \in \mathbb{Q}\}$.

Similarly, \mathbb{R}^n is second countable as well.

\mathbb{R}^ω under the product topology is second countable as well.

$$\left\{ \prod_{n \in \mathbb{N}} U_n : U_n = (a,b) \text{ for } a,b \in \mathbb{Q} \text{ for finitely many } n \text{ and } \mathbb{R} \text{ otherwise} \right\}$$

↳ Why is this set countable?

→ Show that \mathbb{R}^ω under the uniform topology is not second countable.

Hint: Show that the subspace containing sequences of 0s and 1s has the discrete topology so does not have a countable basis. Use this to prove the required.

Theo. (4.3) A subspace of a first (resp. second) countable space is first (second) countable and a countable product of first (resp. second) countable spaces is first (second) countable.

Proof We prove the above for the second countable case. If X has a countable basis \mathcal{B} , then $\{B \cap A : B \in \mathcal{B}\}$ is a countable basis for $A \subseteq X$ as a subspace.

If X_i has countable basis \mathcal{B}_i , then

$$\left\{ \prod U_i : U_i \in \mathcal{B}_i \text{ for finitely many } i \text{ and } U_i = X_i \text{ otherwise} \right\}$$

forms a countable basis of $\prod X_i$. □

Recall that a subset A of a space X is dense in X iff $\bar{A} = X$.

Theo Let X be a second countable space.

- (4.4) (a) Every open covering of X contains a countable subcover.
(b) X has a countable dense subset.

Proof. (a) Let \mathcal{A} be an open covering of X . Let $\mathcal{B} = \{B_n : n \in \mathbb{N}\}$ be a countable basis of X . For each n , if possible, choose an $A_n \in \mathcal{A}$ such that $B_n \subseteq A_n$. Let \mathcal{A}' be the collection of these A_n . It is clearly countable. Further, it covers X . Given $x \in X$, let $B_n \ni x$ be a basis element. Then $x \in A_n$, proving the claim.

(b) For each B_n , let $x_n \in B_n$. Let $D = \{x_n : n \in \mathbb{N}\}$. Then D is a countable dense subset since any basis element of X (and so any open set) intersects D . \square

Def. A space for which every open cover contains a countable subcover is called a **Lindelöf space**.

Lindelöf space

A space having a countable dense subset is said to be **separable**.

Separable space

Lecture 23 - 24/03/21 Separation Axioms

Obviously, any compact set is Lindelöf.

→ Show that \mathbb{R}_l satisfies all countability axioms except the second.

The product of two Lindelöf spaces need not be Lindelöf.

→ Show that \mathbb{R}_l is Lindelöf but $\mathbb{R}_l \times \mathbb{R}_l$ is not.

Hint: Consider $\mathbb{R}_l^2 \setminus \{(x, -x) : x \in \mathbb{R}_l\}$.

A subspace of a Lindelöf space need not be Lindelöf either.

→ Consider the ordered square $I_0^2 = [0, 1]^2$ (under the dictionary order). Show that I_0^2 is compact (and Lindelöf) but the subspace $A = I_0 \times (0, 1)$ is not Lindelöf.

Def. Suppose that one point sets are closed in X . Then X is said to be **regular** if for each $x \in X$ and closed B disjoint from x , there exist disjoint open sets containing x and B .

Regular

Normal

The space is said to be **normal** if for disjoint closed A, B , there exist disjoint open sets containing A and B .

Observe that any normal space is regular and any regular space is Hausdorff.

Lemma. Let one-point sets be closed in X .

(4.5) a) X is regular iff given $x \in X$ and a nbd. U of x , there exists a neighbourhood V of x such that $\bar{V} \subseteq U$.

b) X is normal iff given a closed $A \subseteq X$ and open $U \supseteq A$, there is an open set $V \supseteq A$ with $\bar{V} \subseteq U$.

Proof

a) (Forward) Let $B = X \setminus U$ be closed. There exist disjoint open V and W containing x and B . Then $\bar{V} \cap B = \emptyset$. Therefore, $\bar{V} \subseteq U$.

(Backward) Let $x \in X$ and B disjoint from $\{x\}$ be closed. Let $U = X \setminus B$. Let V be a nbd. of x such that $\bar{V} \subseteq U$. Then V and $X \setminus \bar{V}$ are disjoint open sets containing x and B respectively. Therefore, X is regular.

The argument for (b) is nearly identical, taking A instead of $\{x\}$. \square

Theo. A subspace of a regular space is regular. A product of regular spaces is (4.6) regular.

Proof. Let X be regular and $Y \subseteq X$. One-point sets are closed in Y . Let $x \in Y$ and B a closed subset of Y disjoint from $\{x\}$. Then $\bar{B} \cap Y = B$. Therefore, $x \notin \bar{B}$. Using regularity, let U, V be disjoint open sets containing x and \bar{B} . Then $(U \cap Y)$ and $(V \cap Y)$ are disjoint open sets of Y containing x and B .

Let (X_α) be a family of regular spaces and $X = \prod_\alpha X_\alpha$.

X is Hausdorff, so singletons are closed in X . Let $x \in X$ and U be a neighbourhood of x . Let $\prod_\alpha U_\alpha$ be a basis element containing x and contained in U . For each α , let V_α be a nbd. of x_α such that $\overline{V_\alpha} \subseteq U_\alpha$. If $U_\alpha = X_\alpha$, choose $V_\alpha = X_\alpha$. Then $V = \prod_\alpha V_\alpha$ is a neighbourhood of x in X . Since $\overline{V} = \prod_\alpha \overline{V_\alpha}$, $x \in \overline{V} \subseteq U$, so x is regular. \square

(by Theo 2.9)

→ Show that \mathbb{R}_K is Hausdorff but not regular

Hint: Consider 0 and K .

→ Show that \mathbb{R}_1 is normal.

(4.7)
Theo.

Any second countable regular space is normal

Proof

Let X be regular with countable basis \mathcal{B} . Let A and B be disjoint closed subsets of X . Each $x \in A$ has a nbd. U disjoint from B . Choose a nbd. V of x such that $\overline{V} \subseteq U$. Then, choose an element of \mathcal{B} containing x and contained in V .

This gives a countable covering (U_n) of A by open sets whose closures do not intersect B . Similarly, choose a countable covering (V_n) of B .

$\bigcup U_n$ and $\bigcup V_n$ are open sets containing A and B , but need not be disjoint.

For each n , let

$$U'_n = U_n \setminus \bigcup_{i=1}^n \overline{V}_i \quad \text{and} \quad V'_n = V_n \setminus \bigcup_{i=1}^n \overline{U}_i.$$

Each U'_i and V'_i is open. Also, (U'_n) covers A because for any $x \in A$, $x \in U_n$ for some n but $x \notin \overline{V}_i$ for $1 \leq i \leq n$.

The open sets

$$U' = \bigcup U'_n \quad \text{and} \quad V' = \bigcup V'_n$$

are disjoint. It is easy to show that U' and V' are disjoint. \square

(4.8)

Theo. Any metrizable space is normal.

Proof.

Let X be metrizable with metric d . Let A, B be disjoint closed subsets of X . For each $a \in A$, choose ε_a such that $B_d(a, \varepsilon_a) \cap B = \emptyset$.

Choose ε_b similarly. Then, let

$$U = \bigcup_{a \in A} B_d(a, \varepsilon_a/2) \quad \text{and} \quad V = \bigcup_{b \in B} B_d(b, \varepsilon_b/2).$$

It is easy to show that these are disjoint (they are clearly open and contain A, B), completing the proof. \square

Theo. (4.9) Any compact Hausdorff space is normal.

Left as exercise.

Lecture 24 - 26/03/21 Urysohn Lemma and Completely Regular Spaces

Theo. [Urysohn Lemma]

(4.10) Let A, B be disjoint closed subsets of X . If X is normal, then for a closed interval $[a, b]$ in the real line, there exists a continuous map $f: X \rightarrow [a, b]$ such that $f(x) = a$ for every $x \in A$ and $f(x) = b$ for every $x \in B$.

Proof Clearly, it suffices to take $[a, b] = [0, 1]$.

Let $P = \mathbb{Q} \cap [0, 1]$. For each $p \in P$, we define open U_p such that if $p < q$, $\bar{U}_p \subseteq U_q$.

Arrange P as an infinite sequence (p_n) and for convenience, let $p_1 = 0$ and $p_2 = 1$. Let $U_1 = X \setminus B$. Because A is closed and $A \subseteq U_1$, we may choose (by normality) an open U_0 such that $A \subseteq U_0 \subseteq \bar{U}_0 \subseteq U_1$.

In general, let $P_n = \{p_k : 1 \leq k \leq n\}$ and suppose open U_p is defined for $p \in P_n$ such that $p < q \Rightarrow \bar{U}_p \subseteq U_q$.
($n \geq 2$)

Let $r = p_{n+1}$. Since P_{n+1} is finite, it has a simple ordering $<$ (derived from the usual ordering).

Let p_i and p_j be the immediate predecessor and successor respectively in P_{n+1} (Why do these exist?)

Now, choose U_r as an open set such that $\bar{U}_{p_i} \subseteq U_r \subseteq \bar{U}_r \subseteq U_{p_j}$ — such a U_r exists by using normality on the sets \bar{U}_{p_i} and $X \setminus U_{p_j}$.

→ This defines U_p for $p \in P$ such that $p < q \Rightarrow \bar{U}_p \subseteq U_q$.

Extend this to define U_p for all $p \in \mathbb{Q}$ as $U_p = \emptyset$ if $p < 0$ and $U_p = X$ if $p > 1$.

Given $x \in X$, let $\mathcal{Q}(x) = \{p \in \mathbb{Q} : x \in U_p\}$.

Observe that $\mathcal{Q}(x)$ is bounded below (by, say, -1)

→ Let $f(x) = \inf \mathcal{Q}(x) = \inf \{p \in \mathbb{Q} : x \in U_p\}$.

We claim that f is the desired function.

Note that $f(x) \in [0, 1]$ for any $x \in X$ (Why?). (*)

For any $x \in A \subseteq U_0$, $f(x) = 0$.

For any $x \in B$, $x \notin U_p$ for any $p \leq 1$ ($U_1 = X \setminus B$). By (*), $f(x) = 1$.

It remains to show that f is continuous.

Observe that

• if $x \in \bar{U}_r$, $f(x) \leq r$

• if $x \notin U_r$, $f(x) \geq r$

This follows from the denseness of rationals.

Let $x_0 \in X$ and (c, d) be in \mathbb{R} containing $f(x_0)$.

Choose rationals p, q such that $c < p < f(x_0) < q < d$.

Let $U = U_q \setminus \bar{U}_p$.

Then,

• $f(x_0) < q \Rightarrow x_0 \in U_q$
• $f(x_0) > p \Rightarrow x_0 \notin \bar{U}_p$ } $\Rightarrow x_0 \in U$

• Let $x \in U$. Then $f(x) \in U_q \subseteq \bar{U}_q \Rightarrow f(x) \leq q < d$

$f(x) \notin \bar{U}_p \supseteq U_p \Rightarrow f(x) \geq p > c$

$\Rightarrow f(x_0) \in U \subseteq (c, d)$, so f is continuous by Theo 2.1 (iv). \square

↓
open

Observe that the converse holds too — we may take $U = f^{-1}([0, 1/2])$ and $V = f^{-1}((1/2, 1])$

Def A space X is **completely regular** if one-point sets are closed in X and for each $x_0 \in X$ and closed $A \not\ni x_0$, there is a continuous function $f: X \rightarrow [0, 1]$ such that $f(x_0) = 1$ and $f(A) = \{0\}$.

By the Urysohn Lemma, normality implies complete regularity

The axioms are labelled as

T_1 : for any x, y , there are open U, V such that $x \in U \not\subseteq y$ and $y \in V \not\subseteq x$.

T_2 : Hausdorff

T_3 : Regular

$T_{3\frac{1}{2}}$: Completely regular

T_4 : Normal

Theo. A subspace of a completely regular space is completely regular.

(4.11) A product of completely regular spaces is completely regular.

(Go to next page)

Theo. [Urysohn Metrization Theorem]

(4.12) A regular second countable space X is metrizable

Proof. We shall embed X in a metrizable space Y

Let $Y = \mathbb{R}^\omega$ under the product topology. We have already seen that Y is metrizable under the metric

$$D(x, y) = \sup_i \left\{ \frac{\min\{|x_i - y_i|, 1\}}{i} \right\}.$$

(The proof can also be carried out by taking Y as \mathbb{R}^ω under the uniform topology)

We will in fact embed X in $[0, 1]^\omega$.

→ Claim 1 There exists a countable collection of continuous functions $f_n: X \rightarrow [0, 1]$ such that for any $x_0 \in X$ and nbd. U of x_0 , there is some n such that $f_n(x_0) \neq 0$ and $f(x) = 0$ for $x \in X \setminus U$.

- Let (B_n) be a countable basis for X . For each pair n, m with $\overline{B_n} \subseteq B_m$, use Theo. 3.43 to get a continuous function $g_{n,m}: X \rightarrow [0, 1]$ such that $g_{n,m}(B_n) = \{1\}$ and $g_{n,m}(X \setminus B_m) = \{0\}$.

Then given any $x_0 \in X$ and neighbourhood U of x_0 , we can choose a basis element B_m with $x_0 \in B_m \subseteq U$. By regularity and Lemma 3.38, we can let B_n be a basis element with $x_0 \in B_n \subseteq \overline{B_n} \subseteq B_m$.

$(g_{n,m})$ then satisfies our requirements.

→ For this (f_n) , define $F: X \rightarrow Y$ by

$$F(x) = (f_1(x), f_2(x), f_3(x), \dots)$$

- Because Y has the product topology and each f_n is continuous, F is continuous.

- For $x \neq y$, there is some index n such that $f_n(x) \neq 0$ and $f_n(y) = 0$. So, $F(x) \neq F(y)$ and F is injective.

- We must show that F is a homeomorphism of X to $F(X)$. We have already shown that it is a continuous bijection. Let U be open in X .

Let $z_0 \in F(U)$ and $x_0 \in X$ with $F(x_0) = z_0$. Let N be such that $f_N(x_0) \neq 0$ and $f_N(X \setminus U) = \{0\}$.

Let $V = \pi_N^{-1}((0, \infty)) \subseteq \mathbb{R}^\omega$. Let $W = V \cap F(X)$ be open in $F(X)$. Now,

$\pi_N(z_0) = f_N(x_0) > 0$, so $z_0 \in W$. Further, $W \cap F(X \setminus U) = \emptyset$, so $W \subseteq F(U)$.

Therefore, $F(U)$ is open and F is a homeomorphism. \square

Theo. [Embedding Theorem]

(4.13) Let X be a space in which one-point sets are closed. Let $(f_\alpha)_{\alpha \in J}$ be a family of continuous functions $X \rightarrow \mathbb{R}$ such that for any $x_0 \in X$ and nbd. U of x_0 , there is $\alpha \in J$ such that $f_\alpha(x_0) > 0$ and $f_\alpha(x \setminus U) = \{0\}$. Then $F: X \rightarrow \mathbb{R}^J$ defined by $(F(x))_\alpha = f_\alpha(x)$ is an embedding of X in \mathbb{R}^J .

A family of continuous functions that satisfies the hypothesis of the above theorem is said to separate points from closed sets.

For a space in which one-point sets are closed, this is seen to be equivalent to X being completely regular.

Corollary. A space X is completely regular iff it is homeomorphic to $[0,1]^J$ for (4.14) some J .

Theo. [Tietze Extension Theorem]

Let X be normal and A be closed in X .

(a) Any continuous map $A \rightarrow [a,b] \subseteq \mathbb{R}$ may be extended to a continuous map $X \rightarrow [a,b]$.

(b) Any continuous map $A \rightarrow \mathbb{R}$ may be extended to a continuous $X \rightarrow \mathbb{R}$.

The Tietze Extension Theorem can be used to prove the Urysohn Lemma.
(but its proof uses the Urysohn Lemma)

Theo. [Tychonoff's Theorem]

(4.15) An arbitrary product of compact spaces is compact.

Let $(X_\alpha)_{\alpha \in I}$ be compact and $X = \prod_{\alpha} X_\alpha$.

We first prove a couple of lemmas.

Claim 1. Let X be a set and \mathcal{A} a collection of subsets having the finite intersection property. Then there is a \mathcal{D} such that $\mathcal{A} \subseteq \mathcal{D} \subseteq 2^X$, \mathcal{D} has the finite intersection property, and no \mathcal{F} with $\mathcal{D} \subsetneq \mathcal{F} \subseteq 2^X$ has the finite intersection property.

Proof. We use Zorn's Lemma to prove this.

↳ Given a strictly partially ordered set A in which every simply ordered subset has an upper bound, A has a maximal element.

The strict poset we consider is a set of collections of subsets of X .
Let

$\mathcal{C} = \{ \mathcal{B} \subseteq 2^X : \mathcal{A} \subseteq \mathcal{B} \text{ and } \mathcal{B} \text{ has the finite intersection property} \}$
with the strict partial order strict inclusion \subsetneq .

We want to show that \mathcal{C} has a maximal element \mathcal{D} .

Let $\mathcal{B} \subseteq \mathcal{C}$ be a simply ordered subset. It suffices to show that

$$\mathcal{C} = \bigcup_{\mathcal{B} \in \mathcal{B}} \mathcal{B} \in \mathcal{C}$$

and is an upper bound of \mathcal{B} (which is obvious)

It is clear that $\mathcal{A} \subseteq \mathcal{C}$. Let $C_1, C_2, \dots, C_n \in \mathcal{C}$. For each i , choose $B_i \in \mathcal{B}$ such that $C_i \in B_i$.

$\{B_i : 1 \leq i \leq n\}$ is simply ordered by proper inclusion and is finite, so has a maximal element B_k . Then $C_1, C_2, \dots, C_n \in B_k$. Since B_k has the finite intersection property, $\bigcap_{1 \leq i \leq n} C_i \neq \emptyset$, so \mathcal{C} has the finite intersection property.

Using Zorn's Lemma completes the proof. □

Claim 2. Let X be a set and $\mathcal{D} \subseteq 2^X$ be as defined in the previous claim.

- If B is a finite intersection of elements of \mathcal{D} , $B \in \mathcal{D}$.
- If A is a subset of X that intersects every element of \mathcal{D} , $A \in \mathcal{D}$.

Proof a) Let B equal the intersection of finitely many elements in \mathcal{D} and $\mathcal{E} = \mathcal{D} \cup \{B\}$. We show that \mathcal{E} has the finite intersection property, so $\mathcal{E} = \mathcal{D}$.

Take finitely many elements of \mathcal{E} .

→ If none of them is B , their intersection is clearly nonempty.

→ If B is one of them, we can expand B as a finite intersection to get that the overall intersection is non-empty.

b) Left as exercise (idea similar to a)

We now come to the main proof of Tychonoff's Theorem. Let \mathcal{A} be a collection of subsets of X having the finite intersection property.

We show that $\bigcap_{A \in \mathcal{A}} \bar{A} \neq \emptyset$

By Claim 1, choose $\mathcal{D} \supseteq \mathcal{A}$ as defined.

It suffices to show that $\bigcap_{D \in \mathcal{D}} \bar{D} \neq \emptyset$.

Consider for each $\alpha \in J$

$$\mathcal{D}_\alpha = \{\pi_\alpha(D) \cdot D \in \mathcal{D}\} \subseteq 2^{X_\alpha}$$

Because \mathcal{D} has the finite intersection property, so does \mathcal{D}_α .

By compactness, we may choose for each α , $x_\alpha \in X_\alpha$ such that

$$x_\alpha \in \bigcap_{D_\alpha \in \mathcal{D}_\alpha} \bar{D}_\alpha$$

Let $x = (x_\alpha)_{\alpha \in J} \in X$. If we show that $x \in \bar{D}$ for any $D \in \mathcal{D}$, we are done.

Let $D \in \mathcal{D}$ and U_β be a nbd of $x_\beta \in X_\beta$. Since $x_\beta \in \overline{\pi_\beta(D)}$, we can choose $y \in D$ such that $\pi_\beta(y) \in U_\beta \cap \pi_\beta(D)$.

Then, $y \in \pi_\beta^{-1}(U_\beta) \cap D$.

From (b) of Claim 2, every subbasis element containing x belongs to \mathcal{D} .

By (a) of Claim 2, every basis element containing x belongs to \mathcal{D} and intersects every element of \mathcal{D} . Therefore, $x \in \bar{D}$ for all $D \in \mathcal{D}$.