
STOCHASTIC CALCULUS

Amit Rajaraman

Last updated September 27, 2021

Contents

1	Conditional Expectations and Martingales	2
1.1	Conditional Expectations	2
1.2	Discrete Time Processes	3
1.3	Martingales	4
1.4	Stopping Times	5
1.5	Towards Continuous Time	6
1.6	The Wiener Process	7
1.6.1	Introduction and Definitions	8
1.6.2	Existence and Uniqueness	8
1.6.3	Some Properties	10
2	The Itô Integral and Stochastic Differential Equations	12
2.1	The Itô Integral: Motivations and Definitions	12
2.1.1	An Elementary Definition	12
2.1.2	Towards More Generality	13
2.1.3	Extending the Itô Integral	14
2.1.4	Some Properties	15
2.1.5	Itô Calculus	16
2.2	Stochastic Differential Equations	17

§1. Conditional Expectations and Martingales

1.1. Conditional Expectations

To motivate the general definition of conditional expectation, it is helpful to first work in a discrete setting.

Let (Ω, \mathcal{F}, P) be a probability space and X, Z be random variables such that each of them takes some finite set of values $\{x_1, \dots, x_n\}$ and $\{z_1, \dots, z_m\}$ respectively. What would the conditional expectation be if, say, $Z = z_j$? It is given by

$$\mathbf{E}[X \mid Z = z_j] = \sum x_i \Pr[X = x_i \mid Z = z_j].$$

Denote the above random variable by Y – if $Z(\omega) = z_j$, then $Y(\omega)$ is equal to the above quantity. Note that Y is constant on each Z -atom, and as a result, is measurable with respect to the σ -algebra \mathcal{G} consisting of the 2^m possible unions of Z -atoms. Suppose Y takes value y_j on $\{Z = z_j\}$. Then,

$$\begin{aligned} \int_{\{Z=z_j\}} Y \, dP &= y_j \Pr[Z = z_j] \\ &= \sum_i x_i \Pr[X = x_i \mid Z = z_j] \Pr[Z = z_j] \\ &= \sum_i \Pr[X = x_i, Z = z_j] \\ &= \int_{\{Z=z_j\}} X \, dP. \end{aligned}$$

That is, for any $G \in \mathcal{G}$,

$$\int_G X \, dP = \int_G Y \, dP.$$

This is exactly the motivation behind the general definition.

Definition 1.1. Let (Ω, \mathcal{F}, P) be a probability space and X a random variable with $\mathbf{E}[|X|] < \infty$. Let \mathcal{G} be a sub- σ -algebra of \mathcal{F} . Then, there exists a random variable Y such that

- Y is \mathcal{G} -measurable,
- $\mathbf{E}[|Y|] < \infty$, and
- for every $G \in \mathcal{G}$, $\int_G Y \, dP = \int_G X \, dP$.

Moreover, if Y and \tilde{Y} are two random variables satisfying the above, then $Y = \tilde{Y}$ almost surely. Such a random variable Y is called the *conditional expectation* $\mathbf{E}[X \mid \mathcal{G}]$ of X given \mathcal{G} , and we write $Y = \mathbf{E}[X \mid \mathcal{G}]$ almost surely.

The key insight here is that instead of viewing a random variable as something that takes values, we can view it as a partitioning of the space into equivalence classes, and defining the new random variable based on this partition instead. We do not care about what *values* Z takes, we only care about the probability that Z takes a certain value! Since the conditional expectation is almost surely unique, we mean by “the conditional expectation” any such version of a conditional expectation.

For a random variable Z , $\mathbf{E}[X \mid Z]$ is used to denote $\mathbf{E}[X \mid \sigma(Z)]$.

If $F \in \mathcal{F}$ and \mathcal{G} is a sub- σ -algebra of \mathcal{F} , then we define $\Pr[F \mid \mathcal{G}]$ as $\mathbf{E}[\mathbb{1}_F \mid \mathcal{G}]$.

Lemma 1.1. Let (Ω, \mathcal{F}, P) be a probability space and X a random variable in \mathcal{L}^1 . Let \mathcal{G} and \mathcal{H} be sub- σ -algebras of \mathcal{F} .

- (i) If $Y = \mathbf{E}[X \mid \mathcal{G}]$, then $\mathbf{E}[Y] = \mathbf{E}[X]$.
- (ii) If X is \mathcal{G} -measurable, then $\mathbf{E}[X \mid \mathcal{G}] = X$ almost surely.
- (iii) $\mathbf{E}[a_1 X_1 + a_2 X_2 \mid \mathcal{G}] = a_1 \mathbf{E}[X_1 \mid \mathcal{G}] + a_2 \mathbf{E}[X_2 \mid \mathcal{G}]$ almost surely.
- (iv) If $X \geq 0$, then $\mathbf{E}[X \mid \mathcal{G}] \geq 0$ almost surely.
- (v) If $0 \leq X_n \uparrow X$, then $\mathbf{E}[X_n \mid \mathcal{G}] \uparrow \mathbf{E}[X \mid \mathcal{G}]$ almost surely.
- (vi) If $X_n \geq 0$, then $\mathbf{E}[\liminf X_n \mid \mathcal{G}] \leq \liminf \mathbf{E}[X_n \mid \mathcal{G}]$ almost surely.
- (vii) If $|X_n(\omega)| \leq V(\omega)$ for all n , $\mathbf{E}[V] < \infty$, and $X_n \rightarrow X$ almost surely, then $\mathbf{E}[X_n \mid \mathcal{G}] \rightarrow \mathbf{E}[X \mid \mathcal{G}]$ almost surely.
- (viii) If $c : \mathbb{R} \rightarrow \mathbb{R}$ is convex and $\mathbf{E}[|c(X)|] < \infty$, then $\mathbf{E}[c(X) \mid \mathcal{G}] \geq c(\mathbf{E}[X \mid \mathcal{G}])$. In particular, $\|\mathbf{E}[X \mid \mathcal{G}]\|_p \leq \|X\|_p$ almost surely.
- (ix) If \mathcal{H} is a sub- σ -algebra of \mathcal{G} , then $\mathbf{E}[\mathbf{E}[X \mid \mathcal{G}] \mid \mathcal{H}] = \mathbf{E}[X \mid \mathcal{H}]$ almost surely. We often shorten this as $\mathbf{E}[X \mid \mathcal{G} \mid \mathcal{H}]$.
- (x) If Z is \mathcal{G} -measurable and bounded, then $\mathbf{E}[ZX \mid \mathcal{G}] = Z\mathbf{E}[X \mid \mathcal{G}]$ almost surely. This also holds if $p > 1$, $p^{-1} + q^{-1} = 1$, $X = \mathcal{L}^p(\Omega, \mathcal{F}, P)$, and $Z = \mathcal{L}^q(\Omega, \mathcal{G}, P)$.
- (xi) If \mathcal{H} is independent of $\sigma(\sigma(X), \mathcal{G})$, then $\mathbf{E}[X \mid \sigma(\mathcal{G}, \mathcal{H})] = \mathbf{E}[X \mid \mathcal{G}]$ almost surely. In particular, if X is independent of \mathcal{H} , then $\mathbf{E}[X \mid \mathcal{H}] = \mathbf{E}[X]$ almost surely.

For future purposes, we also recall four types of convergence of random variables, which the reader is hopefully familiar with.

Definition 1.2. Let X be a random variable and (X_n) be a sequence of random variables on the probability space (Ω, \mathcal{F}, P) .

- (a) $X_n \rightarrow X$ *almost surely* if $\Pr[\{\omega \in \Omega : X_n(\omega) \rightarrow X(\omega)\}] = 1$.
- (b) $X_n \rightarrow X$ *in probability* if $\Pr[|X_n - X| > \varepsilon] \rightarrow 0$ for every $\varepsilon > 0$.
- (c) $X_n \rightarrow X$ *in \mathcal{L}^p* if $\|X_n - X\|_p \rightarrow 0$.
- (d) $X_n \rightarrow X$ *in law or in distribution* if $\mathbf{E}[f(X_n)] \rightarrow \mathbf{E}[f(X)]$ for any bounded continuous function f .

We also have that (a),(c) \Rightarrow (b) \Rightarrow (d). Also, if $q \leq p$ and $X_n \rightarrow X$ in \mathcal{L}^p , then $X_n \rightarrow X$ in \mathcal{L}^q .

1.2. Discrete Time Processes

Before we get to stochastic processes, we must build a notion of “time” in a probability space. First, let us do so in a discrete setting.

Definition 1.3. Let (Ω, \mathcal{F}, P) be a probability space. A (discrete time) *filtration* is an increasing sequence (\mathcal{F}_n) of σ -algebras $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \dots \subseteq \mathcal{F}$. The quadruple $(\Omega, \mathcal{F}, (\mathcal{F}_n), P)$ is called a *filtered probability space*.

\mathcal{F}_n is essentially the “set of questions” we can ask at time n . For example, if we are flipping a coin and are at time n , we can ask and get the answer to “Did the coin flips up to time n give more heads than tails?” but we cannot know the answer to “Did the coin flips up to time $N > n$ give more heads than tails?”. We can of course compute probabilities corresponding to the latter case, but we cannot answer it with certainty.

Definition 1.4. Let $(\Omega, \mathcal{F}, (\mathcal{F}_n), P)$ be a filtered probability space. A process (X_n) is called *\mathcal{F}_n -adapted* if X_n is \mathcal{F}_n -measurable for each n , and *\mathcal{F}_n -predictable* if X_n is \mathcal{F}_{n-1} -measurable for each n .

We often refer to the above as just adapted or predictable processes if it is clear from context what the filtration is. In the adapted case, X_n represents something in the past or present, whereas in the predictable case, X_n represents something in the past.

Similarly, we can generate a filtration given a process – this is what we shall use more often.

Definition 1.5. Let (Ω, \mathcal{F}, P) be a probability space and (X_n) be a process. The *filtration generated* by (X_n) is defined by $\mathcal{F}_n^X = \sigma(X_0, \dots, X_n)$ for each n .

Note that the process X_n is \mathcal{F}_n^X -adapted by construction.

Even if some sequences don't explicitly have any relation to the (physical) notion of time, it is often useful to think of them as an adapted stochastic process – for example, a sequence of approximations $\mathbf{E}[X | Y_n]$ to $\mathbf{E}[X | Y]$.

1.3. Martingales

Definition 1.6. A process (X_n) is said to be a \mathcal{F}_n -*martingale* if it is \mathcal{F}_n -adapted and it satisfies $\mathbf{E}[X_n | \mathcal{F}_m] = X_m$ almost surely for every $m \leq n$.

We often refer to the above as just a martingale if it is clear from context what the filtration is.

Observe that the final condition is equivalent to $\mathbf{E}[X_n | \mathcal{F}_{n-1}] = X_{n-1}$.

The name ‘martingale’ comes from a betting strategy. We can think of playing a sequence of games, wherein X_n represents our total winnings after the n th game. In general of course, there need not be any relation between the X_i . In a fair setting however, we should make (and lose) no money on average. If we make money, then it is unfair towards the casino, and if we lose money, then it is unfair towards us (most casinos are unfair towards us). That is, if we have totally won X_m dollars at time m , then our expected winnings at some time point $n > m$ should be $\mathbf{E}[X_n | \sigma\{X_0, \dots, X_m\}] = X_m$ as well!

Martingales turn out to be surprisingly pervasive in probability theory and are seen in several places. To begin with, let us relate martingales a bit more to the above betting strategy, while at the same time giving some intuition as to why martingales are important.

Lemma 1.2 (Doob Decomposition). Let (Ω, \mathcal{F}, P) be a probability space, (\mathcal{F}_n) be a filtration, and (X_n) be \mathcal{F}_n -adapted with $X_n \in \mathcal{L}^1$ for every n . Then, we can write $X_n = X_0 + A_n + M_n$ almost surely, where (A_n) is \mathcal{F}_n -predictable and M_n is a \mathcal{F}_n -martingale with $M_0 = 0$. Further, this decomposition is unique.

This is not too difficult to prove on setting $A_n = \sum_{k=1}^n \mathbf{E}[X_k - X_{k-1} | \mathcal{F}_{k-1}]$.

Definition 1.7. Let (M_n) be a martingale and (A_n) be a predictable process. Then $(A \cdot M)_n = \sum_{k=1}^n A_k(M_k - M_{k-1})$, known as the *martingale transform* of M by A , is a martingale, provided that A_n and $(A \cdot M)_n$ are in \mathcal{L}^1 for all n .

Keeping in line with our gambling analogy, here, M_n would represent our total winnings at time n if we were to stake one dollar in the game and A_k represents the number of dollars we stake at time k . The A_n should be predictable because we place our bet *before* the game occurs.

Let us now consider a particular betting strategy, which seems to go against the intuition built thus far for a martingale. First, choose some $a < b$. We play against a single friend¹ of ours. Whenever our own capital sinks below a , we bet one dollar until it exceeds b dollars. At this point, we stop betting money and wait until our capital sinks below a again.

In each ‘round’ from a to b , we make $(b - a)$ dollars. This seems contradictory! It seems to imply that if we played the game for a long time, we would repeatedly make $(b - a)$ dollars and become rich. However, any betting strategy should be a martingale – what are we missing?

If we face k reversals of fortune, we will make $k(b - a)$ dollars. The only part we have not considered is when we hit a dollars for the $(k + 1)$ th time but have not hit b dollars for the $(k + 1)$ th time yet. Since it must balance out, we must make a significant loss before we reach the next b . That is, the expected loss incurred from the last time we started staking money (when we hit a for the $(k + 1)$ th time) should be equal to $k(b - a)$.

The only logical conclusion in the case where the M_n is bounded is that we can only cross a and b a finite number of times! Otherwise, we could (expect to) make infinite money by playing with the above strategy. That is, the M_n must converge to some random variable.

¹They may not be your friend at the end of the game, however.

Lemma 1.3 (Doob’s Upcrossing Lemma). Let (M_n) be a martingale and denote by $U_n(a, b)$ the number of “upcrossings” of $a < b$ up to time n . Then, $\mathbf{E}[U_n(a, b)] \leq \mathbf{E}[(a - M_n)^+]/(b - a)$.

Proof. This is easily proved as in the previous paragraph. Set $C_0 = 0$ and for $k > 0$, $C_k = \mathbb{1}_{C_{k-1}=1} \mathbb{1}_{M_{k-1} < b} + \mathbb{1}_{C_{k-1}=0} \mathbb{1}_{M_{k-1} \leq a}$. Let $X_n = (C \cdot M)_n$. Then $X_n \geq (b - a)U_n(a, b) - (a - M_n)^+$ (Why?). However, X_n is a martingale, so $\mathbf{E}[X_n] = \mathbf{E}[X_0] = 0$, proving the result. ■

We now come to perhaps the most important result in martingale theory.

Theorem 1.4 (Martingale Convergence Theorem). Let (M_n) be a \mathcal{F}_n -martingale such that $\sup_n \mathbf{E}[|M_n|] < \infty$, $\sup_n \mathbf{E}[(M_n)^+] < \infty$, or $\sup_n \mathbf{E}[(M_n)^-] < \infty$. Then, there exists a \mathcal{F}_∞ -measurable random variable $M_\infty \in \mathcal{L}^1$, where $\mathcal{F}_\infty = \sigma(\mathcal{F}_n : n \in \mathbb{N})$, such that $M_n \rightarrow M_\infty$ almost surely.

Proof. It is not too difficult to show that the three conditions (involving expectation) are equivalent. Let $\omega \in \Omega$. First of all, we claim that that $M_n(\omega)$ cannot have distinct limsup and liminf (so it converges) – if we choose some rational a, b such that $\liminf M_n(\omega) < a < b < \limsup M_n(\omega)$, then there would be infinitely many upcrossings. However,

$$\Pr[\exists a, b \in \mathbb{Q} \text{ such that } M_n \text{ crosses } a, b \text{ i.o.}] \leq \sum_{a, b \in \mathbb{Q}} \Pr[M_n \text{ crosses } a, b \text{ i.o.}] = \sum_{a, b \in \mathbb{Q}} \Pr[U_\infty(a, b) = \infty] = 0,$$

where the last step follows from using the Monotone Convergence Theorem on $\mathbf{E}[U_n(a, b)] \leq (|a| + \sup_n \mathbf{E}[|M_n|])/(b - a) < \infty$, so $\mathbf{E}[U_\infty(a, b)] < \infty$.

Therefore, M_n converges to some M_∞ almost surely. Using Fatou’s Lemma, $\mathbf{E}[M_\infty] \leq \liminf \mathbf{E}[|M_n|] < \infty$, so M_∞ is both almost surely finite and in \mathcal{L}^1 . ■

Similar to martingales, we define the following.

Definition 1.8. A \mathcal{F}_n -adapted process (X_n) is said to be a *supermartingale* if it satisfies $\mathbf{E}[X_n | \mathcal{F}_m] \leq X_m$ almost surely for every $m \leq n$ and a *submartingale* if it satisfies $\mathbf{E}[X_n | \mathcal{F}_m] \geq X_m$ almost surely for every $m \leq n$.

For example, the winnings in casinos usually form a supermartingale – they thrive on the fact that you are expected to lose money.

The analogue of Lemma 1.2 for supermartingales is the following.

Lemma 1.5. Let (Ω, \mathcal{F}, P) be a probability space, (\mathcal{F}_n) be a filtration, and (X_n) be \mathcal{F}_n -adapted with $X_n \in \mathcal{L}^1$ for every n . Then, we can write $X_n = X_0 + A_n + M_n$ almost surely, where (A_n) is a non-increasing predictable process and M_n is a martingale with $M_0 = 0$. Further, this decomposition is unique.

Lemma 1.6. Let M_n be an almost surely non-negative supermartingale and $k > 0$. Then

$$\Pr \left[\sup_n M_n \geq k \right] \leq \frac{\mathbf{E}[M_0]}{k}.$$

1.4. Stopping Times

Definition 1.9. A *stopping time* is a random time $\tau : \Omega \rightarrow \{0, 1, \dots, \infty\}$ such that $\{\omega \in \Omega : \tau(\omega) \leq n\} \in \mathcal{F}_n$ for each n .

Continuing with the gambling analogy, this essentially represents the time we leave the table. For example, quitting while we’re ahead and letting $\tau = \inf\{k : X_k \geq 20\}$ is a stopping time. The definition of a stopping time just says that our decision to leave is based solely on whatever events have transpired thus far. As another example, one might decide to leave when we are done with our final upcrossing (as defined before Lemma 1.3). However, it is impossible to determine whether or not this is the case since we cannot look into the future to see if there are no more upcrossings.

Definition 1.10. Let (X_n) be a stochastic process and $\tau < \infty$ be a stopping time. Then X_τ denotes the random variable $X_{\tau(\omega)}(\omega)$, that is, the process X_n evaluated at τ . In general, the stochastic process $X'_n(\omega) = X_{\min\{n, \tau(\omega)\}}(\omega)$ is called the *stopped process*. That is, $X'_n = X_n$ for $n < \tau$ and $X'_n = X_\tau$ for $n \geq \tau$.

As mentioned, this represents our winnings if we stop playing the game at τ .

Lemma 1.7 (Optional Stopping Theorem). Let M_n be a martingale and $\tau < \infty$ be a stopping time. Then $\mathbf{E}[M_\tau] = \mathbf{E}[M_0]$ if any of the following holds

- (a) $\tau < K$ almost surely for some $K \in \mathbb{N}$.
- (b) $|M_n| < \infty$ almost surely for all n .
- (c) $|M_n - M_{n-1}| < \infty$ almost surely for all n and $\mathbf{E}[\tau] < \infty$.

If any of these conditions hold for a supermartingale, then $\mathbf{E}[M_\tau] \leq \mathbf{E}[M_0]$.

Let us now look at the gambling idea we gave earlier of “quitting while we are ahead”. Suppose ξ_1, ξ_2, \dots are iid and each take ± 1 with probability $1/2$ each. Let $M_n = M_0 + \sum_{i=1}^n \xi_i$. It is easy to see that (M_n) forms a martingale. In the gambling context, this just means that at each time step, we gain or lose a dollar with probability $1/2$.

- First of all, note that M_n does *not* converge (Why?). To solidify the notion of quitting while ahead, consider the stopping time $\tau = \inf\{n : M_n \geq 2M_0\}$. We wait until this happens, and then stop playing. The first question we ask is: is $\tau < \infty$? Perhaps unintuitively, the answer is yes. Indeed, $M_{\min\{n, \tau\}} \leq 2M_0$ almost surely for all n . Therefore, we may apply the martingale convergence theorem to get that $M_{\min\{n, \tau\}}$ converges. Further, since M_n only takes integer values and changes by 1 at each step, it is not too difficult to show that convergence can only occur if it “gets stuck” at M_0 , that is, $\tau < \infty$ almost surely. This seems to be a contradiction – a way to earn free money!
- Next, note that while $M_{\min\{n, \tau\}} \rightarrow 2M_0$ almost surely, it does not converge in \mathcal{L}^1 . Indeed, the latter would imply that the expectations converge as well, but this is clearly not the case since $\mathbf{E}[M_{\min\{n, \tau\}}]$ does not converge to $\mathbf{E}[M_\tau]$. Intuitively, a process does not converge in \mathcal{L}^1 only when the “outliers” of the process grow very rapidly. In the current context, this would mean that while we eventually double our capital, we incur massive losses in the middle in order to keep the game fair.
- To quantify this, suppose that there is also some maximum debt amount $(-R)$ we can incur, so the stopping time is now $\kappa = \inf\{n : M_n \geq 2M_0 \text{ or } M_n \leq -R\}$. Since $|M_{\max\{n, \kappa\}}| \leq \max\{R, 2M_0\}$, we can use Lemma 1.7 to get that $\mathbf{E}[M_\kappa] = \mathbf{E}[M_0]$. However, M_κ takes either $-R$ or $2M_0$, so we can actually calculate the probability that each occurs. The conclusion is that in a fair game, no money can be made on average *unless* we allow ourselves to play for an arbitrarily long amount of time and go arbitrarily far into debt.

1.5. Towards Continuous Time

In the continuous setting, we usually work with stochastic processes in either a time period $[0, T]$ or $[0, \infty)$. In either case, a *stochastic process* on the probability space (Ω, \mathcal{F}, P) is a family of (measurable) random variables (X_t) indexed by time t .

Definition 1.11. Let X_t and Y_t be two stochastic processes. Then X_t and Y_t are said to be *indistinguishable* if $\Pr[X_t = Y_t \text{ for all } t] = 1$ and *modifications* of each other if $\Pr[X_t = Y_t] = 1$ for all t .

First of all, it should be mentioned that we are being slightly messy here. The event “ $X_t = Y_t$ for all t ” is equal to $\bigcap_t \{\omega : X_t(\omega) = Y_t(\omega)\}$, which need not even be an element of the σ -algebra! We shall assume, however, that the σ -algebra is sufficiently rich and this set is an element.

Some texts define indistinguishability slightly different and instead say that this set must contain a subset of measure 1. That is, there exists some set A such that $\Pr[A] = 1$ and for every $\omega \in A$ and t , $X_t(\omega) = Y_t(\omega)$.

Second, it should be noted that indistinguishability of two stochastic processes implies that they are modifications of each other (Why?). Further, the converse holds in the discrete setting (countably many time points) since

$$\Pr \left[\bigcap_n \{\omega : X_n(\omega) = Y_n(\omega)\} \right] \geq 1 - \sum_n \Pr [\{\omega : X_n(\omega) \neq Y_n(\omega)\}].$$

The fact that the converse does not hold in general is shown by considering the probability space with $\Omega = [0, \infty)$ and $\mathcal{B}(\Omega)$, the time period as $[0, \infty)$ and the processes as

$$X_t(\omega) = \begin{cases} 1, & t = \omega, \\ 0, & \text{otherwise,} \end{cases} \quad Y_t(\omega) = 0$$

for all $t \geq 0$ and $\omega \in \Omega$.

In the continuous setting, a filtration is given by a family of σ -algebras (\mathcal{F}_t) indexed by time such that $\mathcal{F}_s \subseteq \mathcal{F}_t$ for all $s \leq t$.

Definition 1.12. Let X_t be a stochastic process on some filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$, and time set $\mathbb{T} \subseteq [0, \infty)$ (of the form $[0, T]$ or $[0, \infty)$). Then X_t is said to be

- *adapted* if X_t is \mathcal{F}_t -measurable for all t ,
- *measurable* if the random variable $X : \mathbb{T} \times \Omega \rightarrow \mathbb{R}$ is $\mathcal{B}(\mathbb{T}) \times \mathcal{F}$ -measurable, and
- *progressively measurable* if $X : [0, t] \cap \mathbb{T} \times \Omega \rightarrow \mathbb{R}$ is $\mathcal{B}([0, t] \cap \mathbb{T}) \times \mathcal{F}$ -measurable for all t .

The first definition is familiar to the reader, and the second and third essentially say that

$$Y_t = \int_0^t X_s \, ds$$

is well-defined and \mathcal{F} -measurable (for measurable) or \mathcal{F}_t -measurable (for progressively measurable). In particular, progressive measurability ensures that Y_t is adapted.

There exists a result that says that every adapted measurable process has a modification that is progressively measurable, but we shall not require it.

Life is made much easier if we restrict ourselves to continuous paths, that is, the function $t \mapsto X_t(\omega)$ is continuous for every ω . Indeed, if they are continuous, then it suffices to compare them at a countable dense subset, say the rationals.

Lemma 1.8. Let X_t and Y_t be stochastic processes with continuous paths. If X_t and Y_t are modifications, then they are indistinguishable. If X_t is adapted and measurable, then it is progressively measurable.

Proof. The first result is direct on considering a countably dense subset (say the rational time points) and using the second remark after Definition 1.11.

Construct a sequence of approximate processes $X_k : [0, t] \times \Omega \rightarrow \mathbb{R}$ such that $X_t^k(\omega) = X_t(\omega)$ for all $\omega \in \Omega$ and $t = 0, 2^{-k}, \dots, 2^{-k} \lfloor 2^{kt} \rfloor$ such that the sample paths are piecewise linear. Then for $s \in [0, t]$ and $\omega \in \Omega$, $X_s^k(\omega) \rightarrow X_s(\omega)$ as $k \rightarrow \infty$. Each X_k is $\mathcal{B}([0, t]) \times \mathcal{F}$ -measurable. Since the limit of a sequence of measurable maps is measurable, the result follows. ■

Theorem 1.9 (Martingale Convergence Theorem). Let M_t be a martingale and suppose that each M_t has continuous sample paths. If $\sup_t \mathbf{E}[|M_t|] < \infty$, $\sup_t \mathbf{E}[(M_t)^+] < \infty$, or $\sup_t \mathbf{E}[(M_t)^-] < \infty$, then there exists a \mathcal{F}_∞ -measurable random variable $M_\infty \in \mathcal{L}^1$ such that $M_t \rightarrow M_\infty$ almost surely.

1.6. The Wiener Process

Brownian motion, which is the main focus of this section, can be thought of as the limit of a random walk such that the time step and the mean square displacement both go to 0. We wish to show that this is in fact well-defined.

1.6.1. Introduction and Definitions

Let ξ_n be iid random variables each with 0 mean and unit variance and define

$$x_t(N) = \sum_{n=1}^{\lfloor Nt \rfloor} \frac{\xi_n}{\sqrt{N}}. \quad (1.1)$$

The Wiener process is just some suitable limit of this sequence. The mere existence of a stochastic process which is the limit of the above, as we have remarked already, is not obvious and we shall construct this process explicitly. The basic tool which comes to mind to do this is the central limit theorem. Unfortunately, that does not work here because we have an uncountable collection of random variables (since there is a random variable for each $t \in [0, T]$). We *can* fix the limiting distribution at some finite number of time steps $x_{t_1}(N), \dots, x_{t_n}(N)$ however. That is,

Lemma 1.10. For any finite set of times $t_1 < \dots < t_n$ ($n < \infty$), the n -dimensional random variable $(x_{t_1}(N), \dots, x_{t_n}(N))$ converges in law as $N \rightarrow \infty$ to an n -dimensional random variable $(x_{t_1}, \dots, x_{t_n})$ such that $x_{t_1}, x_{t_2} - x_{t_1}, \dots, x_{t_n} - x_{t_{n-1}}$ are each independent Gaussian random variables with 0 mean and variance $t_1, t_2 - t_1, \dots, t_n - t_{n-1}$.

The above is easily proved since the increments $x_{t_k}(N) - x_{t_{k-1}}(N)$ are independent for any N . Before getting to the definition of a Wiener process, we state the following, which justifies why we can think of the Wiener process to be continuous.

Lemma 1.11. Suppose we have constructed some stochastic process x_t whose finite dimensional distributions are those of Lemma 1.10. Then, x_t has a modification \tilde{x}_t such that $t \mapsto \tilde{x}_t$ is continuous.

The proof of the above is quite similar to the proof we give later of the existence of a Wiener process.

Definition 1.13. A stochastic process W_t is called a *Wiener process* if

- the finite dimensional distributions of W_t are those of Lemma 1.10 and
- the sample paths of W_t are continuous.

An \mathbb{R}^n -valued process $W_t = (W_t^1, \dots, W_t^n)$ is called an *n -dimensional Wiener process* if W_t^1, \dots, W_t^n are independent Wiener processes.

1.6.2. Existence and Uniqueness

To show that a Wiener process is well-defined, we must establish existence and some sort of uniqueness. We first show uniqueness.

Lemma 1.12. If W_t and W'_t are Wiener processes, then the $C([0, \infty))$ -valued random variables $W, W' : \Omega \rightarrow C([0, \infty))$ have the same law.²

The reader might be wondering exactly what σ -algebra \mathcal{C} we are taking on $C([0, \infty))$. We have two options:

- For each t , consider the evaluation map $\pi_t : C([0, \infty)) \rightarrow \mathbb{R}$, $\pi_t(x) = x_t$. Then set $\mathcal{C} = \sigma\{\pi_t : t \in [0, \infty)\}$.
- Take the natural topology on $C([0, \infty))$ as the topology of uniform convergence on compact intervals. Then, take \mathcal{C} as the Borel σ -algebra with respect to this topology.

It turns out that these two σ -algebras are the same, so our intention is unambiguous.

²Two random variables having the same law means that they induce the same probability measure on the measurable space.

Proof. First, we must show that W and W' are in fact measurable (and thus random variables). Since W_t is measurable for every t (it is a stochastic process), $W_t^{-1}(A) \in \mathcal{F}$ for any $A \in \mathcal{B}(\mathbb{R})$. Note that $W_t = \pi_t \circ W$. Now, by our earlier remark, $\mathcal{C} = \sigma\{\pi_t^{-1}(A) : A \in \mathcal{B}(\mathbb{R}), t \in [0, \infty)\}$, so

$$W^{-1}(\mathcal{C}) = \sigma\{W_t^{-1}(A) : A \in \mathcal{B}(\mathbb{R}), t \in [0, \infty)\} \subseteq \mathcal{F}.$$

Therefore, W is a $[0, \infty)$ -valued random variable (and so is W').

Next, we must show that the two random variables have the same law, that is, they induce the same probability measure on $(C([0, \infty)), \mathcal{C})$. To do this, we use Dynkin's π -system lemma. Consider the π -system

$$\mathcal{C}_{\text{cyl}} = \{\pi_{t_1}(A_1) \cap \cdots \cap \pi_{t_n}(A_n) : t_1, \dots, t_n \in [0, \infty) \text{ and } A_1, \dots, A_n \in \mathcal{B}(\mathbb{R})\}.$$

Now, by definition, the finite-dimensional distributions of W and W' are equal, so the laws coincide on \mathcal{C}_{cyl} . The required follows on using Dynkin's π -system lemma. \blacksquare

Next, how do we show existence? First of all, note that it would suffice to construct a Wiener process on $[0, 1]$ alone. We can then iterate to get it on the succeeding intervals. That is,

Lemma 1.13. Let $\{W_t : t \in [0, 1]\}$ be a stochastic process on (Ω, \mathcal{F}, P) that satisfies Definition 1.13. Then there exists a stochastic process $\{W'_t : t \in [0, \infty)\}$ on a probability space $(\Omega', \mathcal{F}', P')$ that satisfies Definition 1.13 for all t .

The proof follows by setting $\Omega' = \Omega \times \Omega \times \cdots$, $\mathcal{F}' = \mathcal{F} \times \mathcal{F} \times \cdots$, and $P' = P \times P \times \cdots$ with each Ω carrying iid $\{W_t^n : t \in [0, 1]\}$, then checking that $W'_t = \sum_{k=1}^{\lfloor t \rfloor} W_1^k + W_{t-\lfloor t \rfloor}^{\lfloor t \rfloor+1}$ satisfies the required.

So, to show existence on $[0, 1]$, the basic idea is to define a sequence W_t^n of random walks with continuous sample paths, such that $\sum_n \sup_{t \in [0, 1]} |W_t^n - W_t^{n+1}| < \infty$ almost surely. This would imply that they almost surely uniformly converge to some stochastic process W_t and further, this W_t has continuous sample paths.

If this is the case, the structure of the finite dimensional distributions is then easy to see.

So how do we construct these random walks?

The random walk W_t^n consists of 2^n points with the adjacent ones connected by straight lines. To go from W_t^n to W_t^{n+1} , we insert 2^n more points between the old points. That is, we keep adding detail to make the curve finer and finer. The question is: how do we add these points to make the limiting curve have the required characteristics?

By our construction, the points of $W_{k2^{-n}}^n$ are already as in Lemma 1.10. In particular, $W_{(k+1)2^{-n}}^n - W_{k2^{-n}}^n$ is independent of $W_{k2^{-n}}^n$ for any k . Now, given W_t^n , let

$$Y_0 = W_{k2^{-n}}^{n+1} = W_{k2^{-n}}^n \text{ and } Y_1 = W_{(k+1)2^{-n}}^{n+1} = W_{(k+1)2^{-n}}^n.$$

We wish to choose a $X = W_{(2k+1)2^{-(n+1)}}^{n+1}$ such that $Y_1 - X$ and $X - Y_0$ are Gaussian with mean 0 and variance 2^{-n} , and $Y_1 - X$, $X - Y_0$, and Y_0 are independent. It is not too difficult to check that $(Y_0 + Y_1)/2 + 2^{-(n+1)/2}\xi$ does the job, where ξ is standard normal and independent of Y_0 and Y_1 .

Now, let us make the recursion less explicit. These tent-like interpolations we have performed are known as *Schauder functions*.

For $n = 0, 1, \dots$ and $k = 1, 3, \dots, 2^n - 1$, define the *Haar wavelet* $H_{n,k}(t)$ as

$$H_{0,1}(t) = 1 \text{ and } H_{n,k}(t) = \begin{cases} 2^{(n-1)/2}, & (k-1)2^{-n} < t \leq k2^{-n}, \\ -2^{(n+1)/2}, & k2^{-n} < t \leq (k+1)2^{-n}, \\ 0, & \text{otherwise.} \end{cases}$$

The Schauder functions are then defined as

$$S_{n,k}(t) = \int_0^t H_{n,k}(s) \, ds.$$

The N th random walk is then defined as

$$W_t^N = \sum_{n=0}^N \sum_{k=1,3,\dots,2^n-1} \xi_{n,k} S_{n,k}(t),$$

where the $\xi_{n,k}$ are iid standard normal.

Now, we must show that these converge uniformly to prove the required. We have

$$\begin{aligned} \Pr \left[\sup_{t \in [0,1]} |W_t^n - W_t^{n-1}| > \varepsilon_n \right] &= \Pr \left[\sup_{k=1,3,\dots,2^n-1} |\xi_{n,k}| > 2^{(n+1)/2} \varepsilon_n \right] \\ &\leq \sum_{k=1,3,\dots,2^n-1} \Pr \left[|\xi_{n,k}| > 2^{(n+1)/2} \varepsilon_n \right] \\ &= \sum_{k=1,3,\dots,2^n-1} \Pr \left[|\xi_{0,1}| > 2^{(n+1)/2} \varepsilon_n \right] \\ &= \sum_{k=1,3,\dots,2^n-1} 2 \Pr \left[e^{\xi_{0,1}} > \exp \left(2^{(n+1)/2} \varepsilon_n \right) \right] \\ &\leq \sum_{k=1,3,\dots,2^n-1} 2 \exp(-2^{(n+1)/2} \varepsilon_n) \mathbf{E}[e^{\xi_{0,1}}] \\ &= \exp \left(n \log 2 + 1/2 - 2^{(n+1)/2} \varepsilon_n \right). \end{aligned}$$

Setting $\varepsilon_n = n^{-2}$,

$$\sum_{n=1}^{\infty} \Pr \left[\sup_{t \in [0,1]} |W_t^n - W_t^{n-1}| > n^{-2} \right] < \infty.$$

Using the Borel-Cantelli lemma, we infer that

$$\sup_{t \in [0,1]} |W_t^n - W_t^{n-1}| \leq n^{-2}$$

almost surely for sufficiently large n . Therefore,

$$\sum_{n=1}^{\infty} \sup_{t \in [0,1]} |W_t^n - W_t^{n+1}| < \infty$$

almost surely. Setting the sample paths of a null set to 0 is an indistinguishable change, so the W_t^n converge uniformly to W_t , which is continuous. Finally, we must check that W_t has the correct finite-dimensional distributions. This is equivalent to showing that for any $t > s > r$, $W_t - W_s$ and W_r are independent.

To do this, a result states that it suffices to show that

$$\mathbf{E}[e^{i\alpha W_r + i\beta(W_t - W_s)}] = e^{\alpha^2 r/2 - \beta^2(t-s)/2}.$$

Showing this however, is direct by considering a sequence of dyadic rationals – numbers of the form $k2^{-n}$ that converge to s and t , then using dominated convergence and the continuity of W_t .

1.6.3. Some Properties

Definition 1.14. Let \mathcal{F}_t be a filtration. A Wiener process W_t is said to be a \mathcal{F}_t -Wiener process if W_t is \mathcal{F}_t -adapted and $W_t - W_s$ is independent of \mathcal{F}_s for any $t > s$.

Given a Wiener process \mathcal{F}_t , we also define its *natural filtration* $\mathcal{F}_t^W = \sigma\{W_s : s \leq t\}$.

It is not too difficult to show that a \mathcal{F}_t -Wiener process is a \mathcal{F}_t -martingale.

Definition 1.15. A \mathcal{F}_t -adapted process X_t is called a \mathcal{F}_t -Markov process if $\mathbf{E}[f(X_t) | \mathcal{F}_s] = \mathbf{E}[f(X_t) | X_s]$ for all $t \geq s$ and all bounded measurable functions f .

Lemma 1.14. A \mathcal{F}_t -Wiener process is a \mathcal{F}_t -Markov process.

Intuitively, the sample paths of the Wiener process should be very irregular due to the randomness. This is stated better as:

Lemma 1.15. With unit probability, the sample paths of a Wiener process are non-differentiable at any rational time t .

Proof. Suppose that W_t is differentiable at some t . Then for sufficiently small h , $(W_{t+h} - W_t)/h < M$ for some finite M . This implies that $\sup_{n \geq 1} n|W_{t+n^{-1}} - W_t| < \infty$. Now,

$$\begin{aligned} \Pr[\sup_{n \geq 1} n|W_{t+n^{-1}} - W_t| < \infty] &\leq \Pr \left[\bigcup_{M \geq 1} \bigcap_{n \geq 1} \{n|W_{t+n^{-1}} - W_t| < M\} \right] \\ &\leq \lim_{M \rightarrow \infty} \inf_{n \geq 1} \Pr[n|W_{t+n^{-1}} - W_t| < M]. \end{aligned}$$

However, $W_{t+n^{-1}} - W_t$ is Gaussian with mean 0 and variance n^{-1} . As a result,

$$\inf_{n \geq 1} \Pr[n|W_{t+n^{-1}} - W_t| < M] = \inf_{n \geq 1} \Pr[|\xi| < Mn^{-1/2}] = 0,$$

where ξ is standard normal. Therefore, W_t is almost surely not differentiable at t . Since the set of rational numbers is countable, the result follows. \blacksquare

For any real-valued function f , define the *total variation* of f on the interval $[a, b]$ by

$$\text{TV}(f, a, b) = \sup_{k \geq 0} \sup_{(t_i) \in P(k, a, b)} \sum_{i=0}^k |f(t_{i+1}) - f(t_i)|,$$

where $P(k, a, b)$ is the set of partitions $t_0 = a < t_1 < \dots < t_{k+1} = b$ of $[a, b]$.

This is essentially the “distance” travelled to go from a to b along the curve.

Lemma 1.16. With unit probability, $\text{TV}(W, a, b) = \infty$ for any $a < b$.

The main goal of this is to give meaning to the stochastic integral, of the form $\int_0^t f_s dW_s$. However, the above makes this highly problematic. Indeed, it may be shown that if a function g is of infinite total variation, there is a continuous function f such that the usual Stieltjes integral of f with respect to g does not exist.

How do we fix this then? The idea in fact arises by considering the total *squared* variation instead. Note that

$$\lim_{n \rightarrow \infty} \sum_{t_i \in \pi_n} (W_{t_{i+1}} - W_{t_i})^2 = b - a$$

almost surely for some sequence of partitions π_n .

Finally, it is interesting to know that the $x_t(N)$ described in Equation (1.1) do in fact converge in law to a Wiener process.

§2. The Itô Integral and Stochastic Differential Equations

2.1. The Itô Integral: Motivations and Definitions

Motivations

For a moment, let us look at a more physical notion – that of white noise. Given a discrete time signal (a_n) , we model the received signal by $x_n = a_n + \xi_n$, where the ξ_n are iid Gaussian random variables with zero mean. This is discrete time white noise, often known as Additive White Gaussian Noise (AWGN).

How would this extend to continuous time? If we associate a standard normal ξ_t for each $t \in \mathbb{R}^{\geq 0}$ such that $\mathbf{E}[\xi_s \xi_t] = 0$ for $s \neq t$, then if we let $\Xi_\varepsilon = \frac{1}{\varepsilon} \int_0^\varepsilon \xi_s ds$, then it is seen that $\mathbf{E}[\Xi_\varepsilon] = \mathbf{Var}[\Xi_\varepsilon] = 0$. This doesn't make any sense, because it would mean that observing the received signal for an arbitrarily small time from 0 would allow us to completely determine what the transmitted signal at time 0 was. This isn't "noise" then, so how do we fix it?

In AWGN, the corrupting noise in any unit time interval is a zero mean Gaussian (with say, unit variance), so maybe we would want that the average white noise in unit time Ξ_1 is standard normal. We also want that ξ_t and ξ_s are independent for $t \neq s$. This just means that $\int \xi_s ds = W_t!$ However, we have seen that W_t is almost surely non-differentiable, so ξ_t is not really a function. That is, a mathematical model for white noise does not exist, at least within this current theory.³

However, fortunately, it turns out that most of the things we want to do work out if we just stick to working with the Wiener process instead of the white noise itself.

Indeed, in the signal analogy, if (a_t) was transmitted, then the received signal would be $x_t = a_t + \xi_t$. This is not meaningful, but integrating on either side, we get

$$X_t = \int_0^t (a_s + \xi_s) ds = \int_0^t a_s ds + W_t.$$

Instead of attempting to estimate a_t from x_t , we could instead try to solve this problem with X_t , fixing our problem.

2.1.1. An Elementary Definition

Henceforth, in this section, fix a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, \infty)}, \mathbb{P})$ and a \mathcal{F}_t -Wiener process W_t . The stochastic integrals are defined with respect to W_t .

Let us now get to a more mathematical formulation of the above handwavy argument. Similar to how we define a general measure-theoretic integral, we begin by defining the Itô integral for a suitable class of "simple" functions. Finally, we aim to define it for the set of stochastic processes that are \mathcal{F}_t adapted. We take the relevant limits in \mathcal{L}^2 .

Let X_t^n be a \mathcal{F}_{t_i} -adaptable random variable in \mathcal{L}^2 that is constant for $t_i \leq t < t_{i+1}$, where t_i for $i = 0, \dots, N+1$ is a finite set of *non-random* jump times, with the convention that $t_0 = 0$ and $t_{N+1} = T$. For this *simple* integrand, we define the stochastic integral

$$I(X^n) = \int_0^T X_t^n dW_t = \sum_{i=0}^N X_{t_i}^n (W_{t_{i+1}} - W_{t_i}).$$

We aim to extend this definition to a more general class of function. To do so, define the following *Itô Isometry*:

$$\mathbf{E} \left[\left(\int_0^T X_t^n dW_t \right)^2 \right] = \sum_{i=0}^N \mathbf{E}[(X_{t_i}^n)^2] (t_{i+1} - t_i) = \mathbf{E} \left[\int_0^T (X_t^n)^2 dt \right] \quad (2.1)$$

Observe that the adaptability is required for the independence of $X_{t_i}^n$ and $W_{t_{i+1}} - W_{t_i}$ and the \mathcal{L}^2 condition is required for $\mathbf{E}[(X_{t_i}^n)^2]$ to be defined.

$I(X^n)$ is a random variable in \mathcal{L}^2 . It is easier to analyze the above equation if we just think of X_n as a measurable

³Certain theories of generalized stochastic processes do lead to a mathematical model for white noise, but we shall not study these.

map from $[0, T] \times \Omega \rightarrow \mathbb{R}$ for now. Let μ_T be the Lebesgue measure on $[0, T]$. Denote by $\|\cdot\|_{2, \mu}$ the \mathcal{L}^2 -norm with respect to the measure μ . Then, (2.1) just says

$$\|I(X_n)\|_{2, \mathbb{P}} = \|X_n\|_{2, \mu_T \times \mathbb{P}}. \quad (2.2)$$

This is precisely why the equation is an isometry – the map $I : \mathcal{L}^2(\mathbb{P}) \rightarrow \mathcal{L}^2(\mu_T \times \mathbb{P})$ preserves \mathcal{L}^2 -distance (at least when applied to adapted integrands)!

Lemma 2.1. Let $X. \in \mathcal{L}^2(\mu_T \times \mathbb{P})$ and suppose there exists a sequence of \mathcal{F}_t -adapted simple processes $X^n \in \mathcal{L}^2(\mu_T \times \mathbb{P})$ such that

$$\|X^n - X.\|_{2, \mu_T \times \mathbb{P}}^2 = \mathbf{E} \left[\int_0^T (X_t^n - X_t)^2 dt \right] \xrightarrow{n \rightarrow \infty} 0. \quad (2.3)$$

Then $I(X.)$ can be defined as the limit in $\mathcal{L}^2(\mathbb{P})$ of the simple integrands $I(X^n)$ and further, this definition does not depend on the choice of simple approximations of X^n .

Proof. As $m, n \rightarrow \infty$,

$$\|X^m - X^n\|_{2, \mu_T \times \mathbb{P}} \leq \|X^m - X.\|_{2, \mu_T \times \mathbb{P}} + \|X. - X^n\|_{2, \mu_T \times \mathbb{P}} \rightarrow 0.$$

By (2.2), $\|I(X^m) - I(X^n)\|_{2, \mathbb{P}} \rightarrow 0$ as $m, n \rightarrow \infty$ as well. That is, $I(X^n)$ is a Cauchy sequence in $\mathcal{L}^2(\mathbb{P})$. Let $I(X.)$ be the limit of this sequence.

Now, let Y^n be another \mathcal{F}_t -adapted simple process that satisfies (2.3) and let $I(X^n)$ converge to $I(Y.)$ in $\mathcal{L}^2(\mathbb{P})$. Then,

$$\|I(Y.) - I(X.)\|_{2, \mathbb{P}} \leq \|I(Y.) - I(Y^n)\|_{2, \mathbb{P}} + \|I(Y^n) - I(X^n)\|_{2, \mathbb{P}} + \|I(X^n) - I(X.)\|_{2, \mathbb{P}}.$$

The first and last terms converge to 0 by definition and the above argument, and the second argument can be shown to do the same using (2.2).

Therefore, $I(Y.) = I(X.)$ almost surely, proving the claim. ■

Now, the question is: what functions $X. \in \mathcal{L}^2(\mu_T \times \mathbb{P})$ can be approximated as a sequence of \mathcal{F}_t -adapted simple processes (as in (2.3))?

It turns out that this is the case for any \mathcal{F}_t -adapted process.

Lemma 2.2. Let $X. \in \mathcal{L}^2(\mu_T \times \mathbb{P})$ be \mathcal{F}_t -adapted. Then, there is a sequence of \mathcal{F}_t -adapted simple processes $X^n \in \mathcal{L}^2(\mu_T \times \mathbb{P})$

In the case where $X.$ is bounded and has continuous sample paths, the simple functions defined by $X_t^n = X_{k2^{-n}}$, where $k2^{-n}T \leq t < (k+1)2^{-n}T$ gets the job done – this is not too difficult to show using the Dominated Convergence Theorem.

While it is not very complicated, we omit the rest of the proof of the above.

Definition 2.1 (Elementary Itô Integral). Let X_t be any \mathcal{F}_t -adapted process in $\mathcal{L}^2(\mu_T \times \mathbb{P})$. Then the Itô integral $I(X.)$, defined as the limit in $\mathcal{L}^2(\mathbb{P})$ of simple integrals $I(X^n)$ exists and is unique (is independent of the choice of the X^n).

For example, show that $\int_0^T W_t dW_t = W_T^2 - T$.

2.1.2. Towards More Generality

Now, we aim to extend the Itô integral even further. Indeed, while the set of \mathcal{F}_t -adapted processes in $\mathcal{L}^2(\mu_T \times \mathbb{P})$ may seem very open, it is still quite restrictive. Further, we would like to extend this to processes on $[0, \infty)$.

To do so, we first define the Itô integral as a stochastic process on $[0, T]$ with continuous sample paths, then extend it to a process on $[0, \infty)$ using a neat trick called *localization*. We then extend the integral to a wider class of integrands.

Let X_t^n be a \mathcal{F}_t -adapted simple process in $\mathcal{L}^2(\mu_T \times \mathbb{P})$ with jump times $(t_i)_{i=0}^{N+1}$. For any $t \leq T$, define the simple integral

$$I_t(X^n) = \int_0^t X_s^n dW_s = \int_0^T \mathbb{1}_{s \leq t} X_s^n dW_s = \sum_{i=0}^N X_{t_i}^n (W_{\min\{t_{i+1}, t\}} - W_{\min\{t_i, t\}}).$$

First of all, $I_t(X^n)$ is a \mathcal{F}_t -martingale. This is not too difficult to show, show that for any i and $r < t$,

$$\mathbf{E}[X_{t_i}^n (W_{\min\{t_{i+1}, t\}} - W_{t_i, t}) \mid \mathcal{F}_r] = X_{t_i}^n (W_{\min\{t_{i+1}, r\}} - W_{t_i, r}).$$

Indeed, this makes sense since the discrete time $I_{t_i}(X^n)$ is a martingale transform, and the Wiener process is a martingale.

Lemma 2.3. Let X_t be a \mathcal{F}_t -adapted process in $\mathcal{L}^2(\mu_T \times \mathbb{P})$. Then the Itô integral $I_t(X)$, $t \in [0, T]$, can be chosen to have continuous sample paths.

Now, how do we extend the Itô integral to $[0, \infty)$ as a stochastic process? A straightforward idea is to require that the integrand is in $\mathcal{L}^2(\mu \times \mathbb{P})$ and \mathcal{F}_t -adapted, where μ is the Lebesgue measure on $[0, \infty)$.

However, localization, which is not a very deep idea, stipulates that we don't need this. To define it on $[0, \infty)$, it suffices to define it on every $[0, T]$, that is, $X \in \bigcap_{T < \infty} \mathcal{L}^2(\mu_T \times \mathbb{P})$. The integrand does not need to be square integrable, it just needs to be *locally* square integrable, that is, square integrable when restricted to any $[0, T]$.

However, we do need to ensure that this definition is consistent. That is, we need to check that $I_t(X)$ does not depend on which $T > t$ we choose.

This is easily shown however, by taking two times $s < T$ and observing that a sequence (X_t^n) we choose for T would also work for s because $\mathcal{L}^2(\mu_T \times \mathbb{P}) \subseteq \mathcal{L}^2(\mu_s \times \mathbb{P})$.

2.1.3. Extending the Itô Integral

Before we get to an actual extension, let us work with stopping times for a bit.

Lemma 2.4. Let X_t be a \mathcal{F}_t -adapted process in $\bigcap_{T < \infty} \mathcal{L}^2(\mu_T \times \mathbb{P})$ and τ a \mathcal{F}_t -stopping time. Then $I_{\min\{t, \tau\}}(X) = I_t(X \cdot \mathbb{1}_{\cdot < \tau})$.

This is not too difficult to show using our standard technique of proving the claim for simple processes, then extending it.

The condition we are working with right now is that

$$\mathbf{E} \left[\int_0^T X_t^2 dt \right] < \infty$$

for all $T < \infty$. Suppose instead that we are in a situation wherein

$$\mathbf{E} \left[\int_0^{\tau_n} X_t^2 dt \right] < \infty$$

for some sequence $\tau_n \uparrow \infty$ of \mathcal{F}_t -stopping times. Then (τ_n) is said to be a *localizing sequence* for X_t .

While X_t itself need not be in $\mathcal{L}^2(\mu_T \times \mathbb{P})$ for any T , we *do* have that $X_t \mathbb{1}_{t < \tau_n}$ is in $\bigcap_{T < \infty} \mathcal{L}^2(\mu_T \times \mathbb{P})$.

In view of the above lemma then, it makes sense to define $I_t(X)$ as $I_t(X \cdot \mathbb{1}_{\cdot < \tau_n})$ for some n sufficiently large so that $t \leq \tau_n$.

As before, there are two issues we must deal with:

- Does it matter which (sufficiently large) n we take?
- Does it matter which localizing sequence we choose?

The first issue is easily taken care of since for $m > n$, $I_t(X \cdot \mathbb{1}_{\cdot < \tau_n}) = I_t(X \cdot \mathbb{1}_{\cdot < \tau_m})$. Indeed, since $t < \tau_n \leq \tau_m$, we have that $\mathbb{1}_{\cdot < \tau_m} \mathbb{1}_{\cdot < \tau_n} = \mathbb{1}_{\cdot < \tau_n}$. Then since $I_t(X \cdot \mathbb{1}_{t < \tau_n}) = I_{\min\{t, \tau_n\}}(X \cdot \mathbb{1}_{t < \tau_m})$, the proof is straightforward.

For the second issue, let (τ_n) and (τ'_n) be two localizing sequences. Then letting $\sigma_n = \min\{\tau_n, \tau'_n\}$ and applying the above method, we get that the integral corresponding to this stopping time is equal to each of the integrals corresponding to τ_n and τ'_n , so it does not matter which sequence we choose.

Now, this still isn't very handy. However, there is a very natural set of processes that do have localizing sequences, namely that which consists of functions that are \mathcal{F}_t -adapted and satisfy

$$\int_0^T X_t^2 dt < \infty \text{ almost surely for all } T < \infty.$$

A localizing sequence is then

$$\tau_n = \inf \left\{ t \leq n : \int_0^t X_s^2 ds \geq n \right\}.$$

The condition on the integrand implies that $\tau_n \uparrow \infty$ and for all $n \in \mathbb{N}$,

$$\int_0^{\tau_n} X_t^2 dt \leq n$$

almost surely. We may now state the final definition⁴ of the Itô integral.

Definition 2.2 (Itô Integral). Let X_t be a \mathcal{F}_t -adapted stochastic process with

$$\Pr \left[\int_0^T X_t^2 dt < \infty \right] = 1 \text{ for all } T < \infty.$$

Then the Itô integral

$$I_t(X) = \int_0^t X_s dW_s$$

is uniquely defined, by localization and choice of continuous modification, as a \mathcal{F}_t -adapted stochastic process on $[0, \infty)$ with continuous sample paths.

2.1.4. Some Properties

In this section, we give some properties of the Itô integral.

Lemma 2.5 (Linearity). Let X_t and Y_t be Itô integrable processes, and $\alpha, \beta \in \mathbb{R}$. Then $I_t(\alpha X + \beta Y) = \alpha I_t(X) + \beta I_t(Y)$.

Proof. For the case where X_t and Y_t are simple, it follows by definition. If (σ_n) and (τ_n) are localizing sequences for each of the processes, then $\min\{\sigma_n, \tau_n\}$ is a localizing sequence for both, which allows us to extend to the general case by localizing on this sequence. ■

Next, we give a slight generalization (to the wider class of processes) of Lemma 2.4.

Lemma 2.6. Let X_t be Itô integrable and τ a \mathcal{F}_t -stopping time. Then

$$\int_0^{\min\{t, \tau\}} X_s dW_s = \int_0^t X_s \mathbb{1}_{s < \tau} dW_s.$$

⁴Technically, there is a more general definition that allows integration with respect to a general martingale and not just the Wiener process, but we omit it here.

Proof. If (σ_n) is a localizing sequence for X_t , then we see that it is also a localizing sequence for $X_t \mathbb{1}_{t < \tau}$. For $t < \sigma_n$, $I_{\min\{t, \tau\}}(X \cdot \mathbb{1}_{t < \sigma_n}) = I_{\min\{t, \tau\}}(X \cdot \mathbb{1}_{t < \tau} \mathbb{1}_{t < \sigma_n})$. The result follows by localizing on σ_n . ■

Now, let us extend the Itô isometry to a more general integrand.

Lemma 2.7. Let $X \in \bigcap_{T < \infty} \mathcal{L}^2(\mu_T \times \mathbb{P})$. Then for any $T < \infty$,

$$\mathbf{E} \left[\int_0^T X_t dW_t \right] = 0 \text{ and } \mathbf{E} \left[\left(\int_0^T X_t dW_t \right)^2 \right] = \mathbf{E} \left[\int_0^T X_t^2 dt \right].$$

Further, X is a \mathcal{F}_t -martingale.

Proof. The results clearly hold for simple integrands. In general, note that if $Y_n \rightarrow Y$ in \mathcal{L}^2 , then $\mathbf{E}[Y_n] \rightarrow \mathbf{E}[Y]$ and $\mathbf{E}[Y_n^2] \rightarrow \mathbf{E}[Y^2]$ (because convergence in \mathcal{L}^2 implies convergence in law), and $\mathbf{E}[Y_n | \mathcal{F}] \rightarrow \mathbf{E}[Y | \mathcal{F}]$ in \mathcal{L}^2 (Why?). ■

Unfortunately, the above need not hold for a X in the more general class of processes. In fact, in the general case, X need not even be in $\mathcal{L}^1(\mathbb{P})$, so the expectation need not be defined.

Corollary 2.8. If $X^n \rightarrow X$ in $\mathcal{L}^2(\mu_T \times \mathbb{P})$, then $I_t(X^n) \rightarrow I_t(X)$ in $\mathcal{L}^2(\mathbb{P})$. If the convergence is sufficiently fast, then $I_t(X^n) \rightarrow I_t(X)$ almost surely.

We can instead get a condition more general than X just being a \mathcal{F}_t -martingale.

Definition 2.3. A \mathcal{F}_t -measurable process X_t is called a \mathcal{F}_t -local martingale if there exists a sequence of \mathcal{F}_t -stopping times $\tau_n \uparrow \infty$ such that $X_{\min\{t, \tau_n\}}$ is a martingale for each n . Such a (τ_n) is called a *reducing sequence* for X_t .

Any Itô integral $I_t(X)$ is a local martingale. Indeed, any localizing sequence is a reducing sequence.

2.1.5. Itô Calculus

The definition of the Itô integral is quite hard to work with, so in this section, we give some tools to make it much handier.

We work in a more general multi-dimensional framework. Suppose we have a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, \infty)}, \mathbb{P})$, on which an m -dimensional \mathcal{F}_t -Wiener process $W_t = (W_t^1, \dots, W_t^m)$ is defined.

We look at \mathcal{F}_t -adapted processes of the form

$$X_t^i = X_0^i + \int_0^t F_s^i ds + \sum_{j=1}^m \int_0^t G_s^{ij} dW_s^j,$$

where F_s^i and G_s^{ij} are \mathcal{F}_t -progressively measurable processes that satisfy

$$\int_0^t |F_s^i| ds < \infty \text{ and } \int_0^t (G_s^{ij})^2 ds < \infty \text{ almost surely}$$

for all $t < \infty$ and i, j .

Definition 2.4 (Itô Process). A process $X_t = (X_t^1, \dots, X_t^n)$ satisfying the above is called an *n -dimensional Itô process*. It is also denoted as

$$X_t = X_0 + \int_0^t F_s ds + \int_0^t G_s dW_s. \quad (2.4)$$

The main result of this section is the following.

Theorem 2.9 (Itô's Lemma). Let $u : [0, \infty) \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a function such that $u(t, x)$ is C^1 with respect to t and C^2 with respect to x . Then $u(t, X_t)$ is an Itô process:

$$u(t, X_t) = u(0, X_0) + \sum_{\substack{1 \leq i \leq n \\ 1 \leq k \leq m}} \int_0^t u_i(s, X_s) G_s^{ik} dW_s^k + \int_0^t \left(u'(s, X_s) + \sum_{1 \leq i \leq n} u_i(s, X_s) F_s^i + \frac{1}{2} \sum_{\substack{1 \leq i, j \leq n \\ 1 \leq k \leq m}} u_{ij}(s, X_s) G_s^{ik} G_s^{jk} \right) ds, \quad (2.5)$$

where $u'(t, x) = \partial u(t, x) / \partial t$ and $u_i(t, x) = \partial u(t, x) / \partial x_i$.

The above might seem quite unwieldy and arbitrary, but it is actually just the analogue of the chain rule for the Itô integral. Before moving on, let us rewrite this a little to make it more compact.

An Itô process is often written as

$$dX_t = F_t dt + G_t dW_t.$$

This is just suggestive notation for the integrals involved in (2.4). Sticking with this notation, (2.5) says

$$du(t, X_t) = u'(t, X_t) dt + \partial u(t, X_t) dX_t + \frac{1}{2} \text{Tr} [\partial^2 u(t, X_t) dX_t (dX_t)^*], \quad (2.6)$$

where $\partial u(t, x)$ is the row vector with $u_i(t, x)$, $\partial^2 u(t, x)$ is the matrix with entries $u_{ij}(t, x)$, and $dX_t^i dX_t^j$ can be manipulated as

$$(dW_t^i)^2 = dt \text{ and } (dt)^2 = dW_t^i dt = dW_t^i dW_t^j = 0 \text{ if } i \neq j.$$

(Check that the two equations are equivalent!)

- If we set the G^{ij} to 0, the third term vanishes and we recover the usual (deterministic) chain rule from calculus.
- The third extra term is essentially a *second* order approximation. To explain it in an extremely handwavy fashion, we have $(dt)^2 = 0$ in deterministic functions, which is why that term never crops up. Here however, the Itô integral *does* contribute. The reason for this could perhaps be attributed to the fact that $\mathbf{E}[(W_b - W_a)^2] = b - a$, so dW_t can be thought of something like \sqrt{dt} . As a result, the squared term involved in the second order term does contribute non-trivially.

In the one-dimensional case, (2.6) reads

$$du(t, X_t) = \frac{\partial u}{\partial t}(t, X_t) dt + \frac{\partial u}{\partial x}(t, X_t) dX_t + \frac{1}{2} \frac{\partial^2 u}{\partial x^2}(t, X_t) (dX_t)^2,$$

which corresponds exactly to the handwavy argument given above.

Suppose that X_t^1 and X_t^2 are two one-dimensional Itô processes and consider $u(t, x_1, x_2) = x_1 x_2$. Then $u \in C^2$, so Itô's rule (in differential form) implies that

$$dX_t^1 X_t^2 = X_t^1 dX_t^2 + X_t^2 dX_t^1 + dX_t^1 dX_t^2.$$

Therefore, the class of Itô processes is closed under multiplication and forms an algebra.

2.2. Stochastic Differential Equations

Stochastic differential expressions are usually encountered written in the form

$$dX_t = b(t, X_t) dt + \sigma(t, X_t) dW_t \text{ and } X_0 = x.$$

As before, this is just suggestive notation for the Itô process

$$X_t = x + \int_0^t b(s, X_s) ds + \int_0^t \sigma(s, X_s) dW_s.$$

For example, if W_t is a m -dimensional Wiener process, A is a $n \times n$ matrix, and B is a $m \times n$ matrix, then the n -dimensional equation

$$dX_t = AX_t dt + B dW_t \text{ and } X_0 = x$$

is called a *linear stochastic differential equation*. It (always) has a unique solution

$$X_t = e^{At} x + \int_0^t e^{A(t-s)} B dW_s.$$

Call that a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is *Lipschitz uniformly on s* if $\|g(s, x) - g(s, y)\| \leq K \|x - y\|$ for some constant $K < \infty$ that does not depend on s .

For now, let us restrict ourselves to some bounded time $[0, T]$. Consider a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ on which a m -dimensional Wiener process W_t is defined. Choose X_0 to be a \mathcal{F}_0 -measurable n -dimensional random variable, We seek a solution to

$$X_t = X_0 + \int_0^t b(s, X_s) ds + \int_0^t \sigma(s, X_s) dW_s. \quad (2.7)$$

We now show existence of a solution to a large class of differential equations.

Theorem 2.10. Suppose that

1. $X_0 \in \mathcal{L}^2(\mathbb{P})$,
2. b, σ are Lipschitz continuous uniformly on $[0, T]$, and
3. $\|b(t, 0)\|$ and $\|\sigma(t, 0)\|$ are bounded on $t \in [0, T]$.

There exists a solution X_t to the associated stochastic differential equation (2.7). Moreover, for this solution, X_t , $b(t, X_t)$, and $\sigma(t, X_t)$ are in $\mathcal{L}^2(\mu_T \times \mathbb{P})$ and the solution is unique \mathbb{P} -almost surely.

Proof. For \mathcal{F}_t -adapted $Y \in \mathcal{L}^2(\mu_T \times \mathbb{P})$, consider the map

$$(\mathfrak{P}(Y))_t = X_0 + \int_0^t b(s, Y_s) ds + \int_0^t \sigma(s, Y_s) dW_s.$$

Our aim is to find a \mathcal{F}_t -adapted process $X \in \mathcal{L}^2(\mu_T \times \mathbb{P})$ such that $\mathfrak{P}(X) = X$. We carry out the proof of existence in three parts.

- First, we claim that \mathfrak{P} maps to a \mathcal{F}_t -adapted process in $\mathcal{L}^2(\mu_T \times \mathbb{P})$.

Note that

$$\|b\|(t, x) \leq \|b(t, x) - b(t, 0)\| + \|b(t, 0)\| \leq K \|x\| + K' \leq C(1 + \|x\|), \quad (*)$$

where K , K' , and C are suitably chosen constants. Doing so similarly for σ , we may assume that C is large enough such that both the above and $\|\sigma(t, x)\| \leq C(1 + \|x\|)$ hold. To show that \mathfrak{P} maps to a process in \mathcal{L}^2 , we shall show that each of the terms on the right side of the inequality

$$\|(\mathfrak{P}(Y))_t\|_{2, \mu_T \times \mathbb{P}}^2 \leq \|X_0\|_{2, \mu_T \times \mathbb{P}}^2 + \left\| \int_0^t b(s, Y_s) ds \right\|_{2, \mu_T \times \mathbb{P}}^2 + \left\| \int_0^t \sigma(s, Y_s) dW_s \right\|_{2, \mu_T \times \mathbb{P}}^2$$

is finite.

– The first term is

$$\|X_0\|_{2, \mu_T \times \mathbb{P}}^2 = T^{1/2} \|X_0\|_{2, \mathbb{P}}^2 < \infty$$

by the first assumption.

– The second term is

$$\left\| \int_0^t b(s, Y_s) ds \right\|_{2, \mu_T \times \mathbb{P}}^2 \leq T^2 \|b(s, Y_s) ds\|_{2, \mu_T \times \mathbb{P}}^2 \leq T^2 C^2 \|1 + \|Y\cdot\|_{2, \mu_T \times \mathbb{P}}\|^2 < \infty,$$

where the first inequality follows by using Jensen's inequality to get $(t^{-1} \int_0^t a_s ds)^2 \leq t^{-1} \int_0^t a_s^2 ds$, the second inequality uses (*), and the third inequality follows from the fact that Y is in $\mathcal{L}^2(\mu_T \times \mathbb{P})$.

– The third term is

$$\left\| \int_0^t \sigma(s, Y_s) dW_s \right\|_{2, \mu_T \times \mathbb{P}}^2 \leq T \|\sigma(\cdot, Y)\|_{2, \mu_T \times \mathbb{P}}^2 \leq TC^2 \|1 + \|Y\cdot\|_{2, \mu_T \times \mathbb{P}}\|^2 < \infty,$$

where the first inequality follows on using the Itô isometry and the rest is as in the previous step.

This proves that the mapping is to an element of $\mathcal{L}^2(\mu_T \times \mathbb{P})$. It is clear that $\mathfrak{P}(Y)$ is \mathcal{F}_t -adapted, proving the claim.

• Second, we claim that \mathfrak{P} is continuous.

That is, we want to show that if $\|Y^n - Y\|_{2, \mu_T \times \mathbb{P}} \rightarrow 0$, then $\|\mathfrak{P}(Y^n) - \mathfrak{P}(Y)\|_{2, \mu_T \times \mathbb{P}} \rightarrow 0$. As in the first part, we get

$$\|\mathfrak{P}(Y^n) - \mathfrak{P}(Y)\|_{2, \mu_T \times \mathbb{P}} \leq T \|b(\cdot, Y^n) - b(\cdot, Y)\|_{2, \mu_T \times \mathbb{P}} + \sqrt{T} \|\sigma(\cdot, Y^n) - \sigma(\cdot, Y)\|_{2, \mu_T \times \mathbb{P}} \leq K(T + \sqrt{T}) \|Y^n - Y\|_{2, \mu_T \times \mathbb{P}},$$

where the second inequality follows for a suitably large K from the uniformly Lipschitz condition. The required follows.

• Finally, we show the actual existence using a method known as *Picard iteration*.

Let Y_t^0 be an arbitrary \mathcal{F}_t -adapted process in $\mathcal{L}^2(\mu_T \times \mathbb{P})$ and for each $n \geq 0$, let $Y^{n+1} = \mathfrak{P}(Y^n)$. If we show that the (Y^n) converge, then we are done, since the (unique \mathbb{P} -almost surely) limit would be a fixed point of \mathfrak{P} (Why?).

To do so, it suffices to show that the sequence is Cauchy. Exactly as in the proof of the above claim (note the measures involved in each of the norms!),

$$\|(\mathfrak{P}(Y))_t - (\mathfrak{P}(Z))_t\|_{2, \mathbb{P}} \leq \sqrt{t} \|b(\cdot, Y) - b(\cdot, Z)\|_{2, \mu_t \times \mathbb{P}} + \|\sigma(\cdot, Y) - \sigma(\cdot, Z)\|_{2, \mu_t \times \mathbb{P}} \leq L \|Y - Z\|_{2, \mathbb{P}},$$

where $L = K(\sqrt{T} + 1)$. In general,

$$\begin{aligned} \|(\mathfrak{P}^n(X))_t - (\mathfrak{P}^n(Z))_t\|_{2, \mu_T \times \mathbb{P}}^2 &= \int_0^T \|(\mathfrak{P}^n(X))_t - (\mathfrak{P}^n(Z))_t\|_{2, \mathbb{P}}^2 \\ &\leq L^{2n} \int_0^T \int_0^{t_1} \int_0^{t_2} \cdots \int_0^{t_{n-1}} \|Z - Y\|_{2, \mu_{t_n} \times \mathbb{P}}^2 dt_n dt_{n-1} \cdots dt_1 \\ &\leq \frac{L^{2n} T^n}{n!} \|Z - Y\|_{2, \mu_T \times \mathbb{P}}^2. \end{aligned}$$

In particular,

$$\sum_{n=0}^{\infty} \|\mathfrak{P}^{n+1}(Y^0) - \mathfrak{P}^n(Y^0)\|_{2, \mu_T \times \mathbb{P}} \leq \|\mathfrak{P}(Y^0) - Y^0\|_{2, \mu_T \times \mathbb{P}} \sum_{n=0}^{\infty} \sqrt{\frac{L^{2n} T^n}{n!}} < \infty,$$

completing this part of the proof.

Next, suppose that X is the above obtained solution and Y is another solution. We prove uniqueness in two steps.

- If $Y \in \mathcal{L}^2(\mu_T \times \mathbb{P})$, then for any n ,

$$\|X - Y\|_{2, \mu_T \times \mathbb{P}}^2 = \|\mathfrak{P}^n(X) - \mathfrak{P}^n(Y)\|_{2, \mu_T \times \mathbb{P}}^2 \leq \sqrt{\frac{L^{2n} T^n}{n!}} \|X - Y\|_{\mu_T \times \mathbb{P}}^2.$$

Letting $n \rightarrow \infty$, we see that the expression on the left must be 0, so X and Y are equal $\mu_T \times \mathbb{P}$ -almost surely.

- To complete the proof, we show that any solution Y_t with $Y_0 = X_0$ must be in $\mathcal{L}^2(\mu_T \times \mathbb{P})$.

■